

## Refine Search

### Search Results -

Terms	Documents
L17 WITH inactivated	2

Database:

US Pre-Grant Publication Full-Text Database  
 US Patents Full-Text Database  
 US OCR Full-Text Database  
 EPO Abstracts Database  
 JPO Abstracts Database  
 Derwent World Patents Index  
 IBM Technical Disclosure Bulletins

Search:

L27

Search Forms

Search

Results

User Searches

Preferences

Logout




### Search History

 DATE: Saturday, February 25, 2006    [Printable Copy](#)    [Create Case](#)

<u>Set Name</u> side by side	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u> result set
<i>DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI,TDBD; THES=ASSIGNEE; PLUR=YES; OP=ADJ</i>			
<a href="#">L27</a>	l17 WITH inactivated	2	<a href="#">L27</a>
<a href="#">L26</a>	l17 NEAR inactivated	2	<a href="#">L26</a>
<a href="#">L25</a>	l17 SAME inactivate	0	<a href="#">L25</a>
<a href="#">L24</a>	l17 NEAR inactivate	0	<a href="#">L24</a>
<a href="#">L23</a>	l21 and l17	2	<a href="#">L23</a>
<a href="#">L22</a>	L21 or l21	250	<a href="#">L22</a>
<a href="#">L21</a>	mycobacterium vaccae	250	<a href="#">L21</a>
<a href="#">L20</a>	mycobacterium	20667	<a href="#">L20</a>
<a href="#">L19</a>	mycobaterium	132	<a href="#">L19</a>
<a href="#">L18</a>	mycobacteriaum	4	<a href="#">L18</a>
<a href="#">L17</a>	bacilli calmette-guerin	1487	<a href="#">L17</a>
<a href="#">L16</a>	adjuvant OR adjuvants	122183	<a href="#">L16</a>
<a href="#">L15</a>	L13 and inactivated	2	<a href="#">L15</a>
<a href="#">L14</a>	L13 and inactivate	0	<a href="#">L14</a>
<a href="#">L13</a>	L12 and adjuvant	4	<a href="#">L13</a>

<u>L12</u>	L11 or l10	4	<u>L12</u>
<u>L11</u>	6699483.pn.	2	<u>L11</u>
<u>L10</u>	6972128.pn.	2	<u>L10</u>
<u>L9</u>	inactivated adjuvants	13	<u>L9</u>
<u>L8</u>	L7 and l4	0	<u>L8</u>
<u>L7</u>	L6 and l5	12	<u>L7</u>
<u>L6</u>	bacille calmette-guerin	3108	<u>L6</u>
<u>L5</u>	mycobacterium vaccae	250	<u>L5</u>
<u>L4</u>	inactivate	47072	<u>L4</u>
<u>L3</u>	L2 and liver	2	<u>L3</u>
<u>L2</u>	L1 and bone	2	<u>L2</u>
<u>L1</u>	20050019336	2	<u>L1</u>


END OF SEARCH HISTORY

## Product Description

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's [Material Transfer Agreement](#) or, in certain cases, an MTA specified by the depositing institution.

Customers in Europe, Australia, Hong Kong, India, Japan, Korea, New Zealand, Singapore and Taiwan, R.O.C. must contact a [local distributor](#) for pricing information and to place an order for ATCC cultures and products.

### Cell Biology

<b>ATCC® Number:</b>	<b>CRL-1740™</b>	<b>Price:</b>	<b>\$203.00</b>
	<div>Order this item</div>		
<b>Designations:</b>	LNCaP clone FGC [LNCaP.FGC]	<b>Depositors:</b>	JS Horoszewicz
<b>Biosafety Level:</b>	1	<b>Shipped:</b>	frozen
<b>Medium &amp; Serum:</b>	<u>See Propagation</u>	<b>Growth Properties:</b>	adherent, single cells and loosely attached clusters
<b>Organism:</b>	<i>Homo sapiens</i> (human)	<b>Morphology:</b>	epithelial
			
<b>Source:</b>	<b>Organ:</b> prostate <b>Disease:</b> carcinoma <b>Tumor stage:</b> <b>Derived from metastatic site:</b> left supraclavicular lymph node		
<b>Cellular Products:</b>	human prostatic acid phosphatase; prostate specific antigen [21889]		
<b>Permits/Forms:</b>	In addition to the <u>MTA</u> mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please <u><a href="#">click here</a></u> for information regarding the specific requirements for shipment to your location.		

### Related Cell Culture Products

<b>Restrictions:</b>	Distribution of this material for commercial purposes will require execution of a Non-exclusive License Agreement. At the time of placing an order, customers must send a request to <a href="mailto:licensing@atcc.org">licensing@atcc.org</a> . Orders will be shipped when Customer Service receives confirmation from our Licensing officer.
<b>Isolation:</b>	<b>Isolation date:</b> 1977
<b>Applications:</b>	transfection host ( <a href="#">technology from amaxa</a> )
<b>Receptors:</b>	androgen receptor, positive; estrogen receptor, positive [23045]
<b>Tumorigenic:</b>	Yes, in soft agar Yes, the cells are tumorigenic in nude mice
<b>Cytogenetic Analysis:</b>	This is a hypotetraploid human cell line. The modal chromosome number was 84, occurring in 22% of cells. However, cells with chromosome counts of 86 (20%) and 87 (18%) also occurred at high frequencies. The rate of cells with higher ploidies was 6.0%.
<b>Age:</b>	50 years adult

<b>Gender:</b>	male
<b>Ethnicity:</b>	Caucasian
<b>Comments:</b>	<p>LNCaP clone FGC was isolated in 1977 by J.S. Horoszewicz, et al., from a needle aspiration biopsy of the left supraclavicular lymph node of a 50-year-old Caucasian male (blood type B+) with confirmed diagnosis of metastatic prostate carcinoma. [21889]</p> <p>These cells are responsive to 5-alpha-dihydrotestosterone (growth modulation and acid phosphatase production). [23045]</p> <p>The cells do not produce a uniform monolayer, but grow in clusters which should be broken apart by repeated pipetting when subcultures are prepared.</p> <p>They attach only lightly to the substrate, do not become confluent and rapidly acidify the medium.</p> <p>Growth is very slow.</p> <p>The cells should be allowed to incubate undisturbed for the first 48 hours after subculture.</p> <p>When flask cultures are shipped, the majority of the cells become detached from the flask and float in the medium.</p> <p>Upon receipt, incubate the flask (in the usual position for monolayer cultures) for 24 to 48 hours to allow the cells to re-attach.</p> <p>The medium can then be removed and replaced with fresh medium.</p> <p>If desired, the contents of the flask can be collected, centrifuged at 300 X g for 15 minutes, resuspended in 10 ml of medium and dispensed into a single flask.</p>
<b>Propagation:</b>	<p><b>ATCC complete growth medium:</b> RPMI 1640 medium with 2 mM L-glutamine adjusted to contain 1.5 g/L sodium bicarbonate, 4.5 g/L glucose, 10 mM HEPES, and 1.0 mM sodium pyruvate, 90%; fetal bovine serum, 10%</p> <p><b>Temperature:</b> 37.0C</p> <p><b>Atmosphere:</b> air, 95%; carbon dioxide (CO<sub>2</sub>), 5%</p>
<b>Subculturing:</b>	<p><b>Protocol:</b></p> <ol style="list-style-type: none"> <li>1. Remove and discard culture medium.</li> <li>2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin- 0.53 mM EDTA solution to remove all traces of serum that contains trypsin inhibitor.</li> <li>3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.</li> <li>4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.</li> <li>5. Add appropriate aliquots of the cell suspension to new culture vessels. Maintain cultures at a cell concentration between 1 X 10<sup>4</sup> and 2 X 10<sup>5</sup> cells/cm<sup>2</sup>.</li> <li>6. Incubate cultures at 37°C.</li> </ol> <p><b>Subcultivation ratio:</b> A subcultivation ratio of 1:3 to 1:6 is recommended</p> <p><b>Medium renewal:</b> Twice per week</p>
<b>Preservation:</b>	<p><b>Freeze medium:</b> Complete growth medium supplemented with 5% (v/v) DMSO</p> <p><b>Storage temperature:</b> liquid nitrogen vapor phase</p>
<b>Doubling Time:</b>	about 34 hours
<b>Related Products:</b>	<p>Recommended medium (without the additional supplements or serum described under ATCC Medium): <a href="#">ATCC 30-2001</a></p> <p>recommended serum: <a href="#">ATCC 30-2020</a></p> <p>purified DNA: <a href="#">ATCC CRL-1740D</a></p>
<b>References:</b>	<p>21889: Murphy GP, editor. Models for prostate cancer. 37: New York: Liss; 1980, pp. 115-132.</p> <p>22410: Gibas Z , et al. A high-resolution study of chromosome changes in a human prostatic carcinoma cell line (LNCaP). Cancer Genet. Cytogenet. 11: 399-404, 1984. PubMed: <a href="#">6584201</a></p> <p>23045: Horoszewicz JS , et al. LNCaP model of human prostatic carcinoma. Cancer Res. 43: 1809-1818, 1983. PubMed: <a href="#">6831420</a></p> <p>32283: Hu SX , et al. Development of an adenovirus vector with tetracycline-regulatable</p>

human tumor necrosis factor alpha gene expression. Cancer Res. 57: 3339-3343, 1997. PubMed: [9269991](#)  
33090: Boffa LC , et al. Invasion of the CAG triplet repeats by a complementary peptide nucleic acid inhibits transcription of the androgen receptor and TATA-binding protein genes and correlates with refolding of an active nucleosome containing a unique AR gene sequence. J. Biol. Chem. 271: 13228-13233, 1996. PubMed: [8662737](#)

#### Notices and Disclaimers

ATCC products are intended for laboratory research purposes only. They are not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for nonprofit institutions in the United States and Canada. Cultures that are ordered as test tubes or flasks will carry an additional laboratory fee. Fees for permits, shipping, and handling may apply.

You may continue your ATCC Number search by typing in your search criteria below or returning to the [ATCC Search Page](#). For more information please review the [Search Help](#).

#### **Home Page Archive**

[Home](#) [Ordering Info](#) [Quick Order](#) [Support](#) [About ATCC](#) [Contact Us](#)  
[Privacy Policy](#) [Terms of Use](#) [ATCC MTA](#)

© 2006 American Type Culture Collection (ATCC).  
All rights reserved.



Your Discoveries  
Begin with US.

Search Catalog:

[Home](#) | [Ordering Info](#) | [Quick Order](#) | [Cart](#)

## Product Description

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's [Material Transfer Agreement](#) or, in certain cases, an MTA specified by the depositing institution.

Customers in Europe, Australia, Hong Kong, India, Japan, Korea, New Zealand, Singapore and Taiwan, R.O.C. must contact a [local distributor](#) for pricing information and to place an order for ATCC cultures and products.

### Cell Biology

**ATCC® Number:** HTB-81™ **Price:** \$203.00  
[Order this item](#)

**Designations:** DU 145

**Depositors:** KR Stone

**Biosafety Level:** 1

**Shipped:** frozen

**Medium & Serum:** [See Propagation](#)

**Growth Properties:** adherent

**Organism:** *Homo sapiens* (human)

**Morphology:** epithelial

**Source:** **Organ:** prostate  
**Disease:** carcinoma  
**Tumor stage:**  
**Derived from metastatic site:** brain

**Permits/Forms:** In addition to the [MTA](#) mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

### Related Cell Culture Products

**Isolation:** (DU 145 was isolated by K.R. Stone et al from a lesion in the brain of a patient with metastatic carcinoma of the prostate and a 3 year history of lymphocytic leukemia.)

**Applications:** transfection host ([technology from amaxa](#))

**Tumorigenic:** Yes, in nude mice; forms adenocarcinoma (grade II) consistent with prostatic primary

**Antigen Expression:** Blood Type O; Rh+

**Cytogenetic Analysis:** This is a hypotriploid human cell line. Both 61 and 62 chromosome numbers had the highest rate of occurrence in 30 metaphase counts. The rate of higher ploidies was 3%. The t(11q12q), del(11)(q23), 16q+, del(9)(p11), del(1)(p32) and 6 other marker chromosomes were found in most cells. The N13 was usually absent. The Y chromosome is abnormal through translocation to an unidentified chromosomal segment. The X chromosome was present in single copy.

**Isoenzymes:** AK-1, 1; ES-D, 1; G6PD, B; GLO-I, 2; Me-2, 1-2; PGM1, 1; PGM3, 2

**Age:** 69 years

**Gender:** male

**Ethnicity:** Caucasian

<b>Comments:</b>	The line is not detectably hormone sensitive, is only weakly positive for acid phosphatase and isolated cells form colonies in soft agar. The cells do not express prostate antigen. Ultrastructural analyses of both the cell line and original tumor revealed microvilli, tonofilaments, desmosomes, any mitochondria, well developed Golgi and heterogenous lysosomes.
<b>Propagation:</b>	<b>ATCC complete growth medium:</b> Minimum essential medium (Eagle) with 2 mM L-glutamine and Earle's BSS adjusted to contain 1.5 g/L sodium bicarbonate, 0.1 mM non-essential amino acids, and 1.0 mM sodium pyruvate, 90%; fetal bovine serum, 10% <b>Temperature:</b> 37.0C <b>Atmosphere:</b> air, 95%; carbon dioxide (CO <sub>2</sub> ), 5%
<b>Subculturing:</b>	<b>Protocol:</b> <ol style="list-style-type: none"> <li>1. Remove and discard culture medium.</li> <li>2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin - 0.53 mM EDTA solution to remove all traces of serum which contains trypsin inhibitor.</li> <li>3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37C to facilitate dispersal.</li> <li>4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.</li> <li>5. Add appropriate aliquots of the cell suspension to new culture vessels.</li> <li>6. Incubate cultures at 37C.</li> </ol> <b>Subcultivation ratio:</b> A subcultivation ratio of 1:4 to 1:6 is recommended <b>Medium renewal:</b> 2 to 3 times per week
<b>Preservation:</b>	<b>Freeze medium:</b> Complete growth medium, 95%; DMSO, 5% <b>Storage temperature:</b> liquid nitrogen vapor temperature
<b>Related Products:</b>	Recommended medium (without the additional supplements or serum described under ATCC Medium): ATCC <a href="#">30-2003</a> recommended serum: ATCC <a href="#">30-2020</a> purified DNA: ATCC <a href="#">HTB-81D</a> 0.25% (w/v) Trypsin - 0.53 mM EDTA in Hank' BSS (w/o Ca++, Mg++): ATCC <a href="#">30-2101</a> Cell culture tested DMSO: ATCC <a href="#">4-X</a>
<b>References:</b>	22289: Papsidero LD , et al. Prostate antigen: a marker for human prostate epithelial cells. J. Natl. Cancer Inst. 66: 37-42, 1981. PubMed: <a href="#">6935463</a> 22858: Stone KR , et al. Isolation of a human prostate carcinoma cell line (DU 145). Int. J. Cancer 21: 274-281, 1978. PubMed: <a href="#">631930</a> 23028: Mickey DD , et al. Heterotransplantation of a human prostatic adenocarcinoma cell line in nude mice. Cancer Res. 37: 4049-4058, 1977. PubMed: <a href="#">908039</a> 23226: Pollack MS , et al. HLA-A, B, C and DR alloantigen expression on forty-six cultured human tumor cell lines. J. Natl. Cancer Inst. 66: 1003-1012, 1981. PubMed: <a href="#">7017212</a> 32283: Hu SX , et al. Development of an adenovirus vector with tetracycline-regulatable human tumor necrosis factor alpha gene expression. Cancer Res. 57: 3339-3343, 1997. PubMed: <a href="#">9269991</a> 32341: Sheng S , et al. Maspin acts at the cell membrane to inhibit invasion and motility of mammary and prostatic cancer cells. Proc. Natl. Acad. Sci. USA 93: 11669-11674, 1996. PubMed: <a href="#">8876194</a> 32460: Carter RE , et al. Prostate-specific membrane antigen is a hydrolase with substrate and pharmacologic characteristics of a neuropeptidase. Proc. Natl. Acad. Sci. USA 93: 749-753, 1996. PubMed: <a href="#">8570628</a> 32486: Nupponen NN , et al. Genetic alterations in prostate cancer cell lines detected by comparative genomic hybridization. Cancer Genet. Cytogenet. 101: 53-57, 1998. PubMed: <a href="#">9460501</a> 32768: Robinson D , et al. A tyrosine kinase profile of prostate carcinoma. Proc. Natl. Acad. Sci. USA 93: 5958-5962, 1996. PubMed: <a href="#">8650201</a> 32916: Su ZZ , et al. Surface-epitope masking and expression cloning identifies the human prostate carcinoma tumor antigen gene PCTA-1 a member of the galectin gene family. Proc. Natl. Acad. Sci. USA 93: 7252-7257, 1996. PubMed: <a href="#">8692978</a> 32925: Zhu X , et al. Cell cycle-dependent modulation of telomerase activity in tumor cells. Proc. Natl. Acad. Sci. USA 93: 6091-6095, 1996. PubMed: <a href="#">8650224</a> 33090: Boffa LC , et al. Invasion of the CAG triplet repeats by a complementary peptide

nucleic acid inhibits transcription of the androgen receptor and TATA-binding protein genes and correlates with refolding of an active nucleosome containing a unique AR gene sequence. J. Biol. Chem. 271: 13228-13233, 1996. PubMed: [8662737](#)

#### Notices and Disclaimers

ATCC products are intended for laboratory research purposes only. They are not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for nonprofit institutions in the United States and Canada. Cultures that are ordered as test tubes or flasks will carry an additional laboratory fee. Fees for permits, shipping, and handling may apply.

You may continue your ATCC Number search by typing in your search criteria below or returning to the [ATCC Search Page](#). For more information please review the [Search Help](#).

HTB-81	ATCC Number Search	Clear Search
--------	--------------------	--------------

#### **Home Page Archive**

[Home](#) [Ordering Info](#) [Quick Order](#) [Support](#) [About ATCC](#) [Contact Us](#)  
[Privacy Policy](#) [Terms of Use](#) [ATCC MTA](#)

© 2006 American Type Culture Collection (ATCC).  
All rights reserved.